

Recent results on moments of parton distribution functions *

I. Wetzorke^a, K. Jansen^a, F. Palombi^b, A. Shindler^a

^aNIC/DESY Zeuthen, Platanenallee 6, D-15738 Zeuthen, Germany

^bE. Fermi Research Center, c/o Compendio Viminale, pal. F, I-00184 Rome, Italy

We report on recent results for the pion and nucleon matrix element of the twist-2 operator corresponding to the average momentum of non-singlet quark densities. We discuss finite size effects for the nucleon matrix element and present first preliminary results for the non-perturbative renormalisation from full dynamical simulations.

1. INTRODUCTION

Phenomenological fits to experimental data provide results for moments of parton distribution functions which allow a direct comparison with lattice calculations. In a recent paper [1] we were able to compute a continuum limit value of the lowest moment of a twist-2 operator, corresponding to the average momentum of non-singlet quark densities, in pion states. In obtaining the renormalisation group invariant matrix element, we have controlled important systematic errors, such as non-perturbative renormalisation [2], finite size effects [3] and effects of a non-vanishing lattice spacing. The crucial limitation of our calculation was the use of the quenched approximation.

In the scope of our current investigations is the comparison of finite size effects for pion and nucleon matrix elements. In addition we concentrate on computing non-perturbatively the evolution of the twist-2 operators, using the Schrödinger functional finite-size technique, on $N_f = 2$ dynamical configurations provided by the ALPHA collaboration [5].

2. FINITE SIZE EFFECTS

Finite size effects for nucleon matrix elements have not been studied in detail before, but might influence considerably the precise determination of moments of parton distribution functions from lattice QCD. We have investigated the effects of limited lattice extent at $\beta=6.0$ ($a \simeq 0.093$ fm) with the non-perturbatively improved clover fermion action and lattice sizes $(12 - 32)^3 \times 32$.

The twist-2 non-singlet operator for a flavour

doublet of fermions belongs to two irreducible representations of the lattice $H(4)$ group

$$\begin{aligned}\mathcal{O}_{12}(x) &= \frac{1}{4} \bar{\psi}(x) \gamma_{\{1} \overleftrightarrow{D}_{2\}} \tau^3 \psi(x) \\ \mathcal{O}_{44}(x) &= \frac{1}{2} \bar{\psi}(x) \left[\gamma_4 \overleftrightarrow{D}_4 - \frac{1}{3} \sum_{k=1}^3 \gamma_k \overleftrightarrow{D}_k \right] \tau^3 \psi(x) .\end{aligned}$$

For the study of finite size effects (FSE) we concentrate solely on the $\mathcal{O}_{44}$ operator, since it can be computed at zero external momentum and thus provides a reliable signal. In particular we use Schrödinger functional (SF) boundary conditions [6] for our computation, which allows the extraction of the matrix element from one large plateau around the middle of the time extent T of the lattice. The correlation function of the matrix element f_M is obtained by inserting the $\mathcal{O}_{44}$ operator between two SF nucleon states $\mathbb{S}$ and $\mathbb{S}'$ at the time boundaries $t = 0$ and T and suitable normalisation with the boundary-to-boundary correlator C_1



$$\begin{aligned}f_M(x_0/L) &= -\frac{1}{2} \sum_x \langle \mathbb{S}' | \mathcal{O}_{44}(x) | \mathbb{S} \rangle \\ &\sim e^{-m_N T} \langle N | \mathcal{O}_{44} | N \rangle \{1 + \dots\} \\ C_1 &= -\frac{1}{2} \langle \mathbb{S}' \mathbb{S} \rangle \sim e^{-m_N T} .\end{aligned}$$

The bare nucleon matrix element is defined by

$$\langle x \rangle_{u-d}^{bare} \equiv \frac{2\kappa}{m_N} \langle N | \mathcal{O}_{44} | N \rangle = \frac{2\kappa}{m_N} f_M(x_0/L) / C_1 .$$

In order to obtain the infinite volume limit nucleon mass and matrix element we performed a purely phenomenological fit $X(L) = c_0 + c_1/L^{3/2} \exp(-c_2 L)$, where $X = m_N a$ or $\langle x \rangle_{u-d}$. Please note that a power-law fit ansatz would describe the data almost equally well. In figs. 1 and

*Talk presented by I. Wetzorke. The work was supported by the DFG under SFB/TR 09-03.

Figure 1. Finite size effects for $m_N a$ Figure 2. Finite size effects for nucleon $\langle x \rangle_{u-d}$

2 we show the relative systematic deviation from the infinite volume limit values for both the nucleon mass and matrix element. While the nucleon mass shows a deviation of about 4% at $m_\pi L \simeq 5$, the nucleon matrix element is overestimated by 10-30% at this value of $m_\pi L$. In both cases one can be sure to exclude FSE only for $m_\pi L > 7$. For the pion matrix element we observed a FSE of about 3% at $m_\pi L \simeq 5$, while the pion mass is found to be free from FSE already for $m_\pi L > 4$.

These observations lead to the expectation that the strong FSE for nucleon matrix elements could be one ingredient for the deviation of current lattice QCD estimates of $\langle x \rangle_{u-d}$ in comparison with experimental data.

3. $N_f = 2$ CONTINUUM STEP SCALING FUNCTION

The strategy used to compute the non-perturbative evolution of the operators $\mathcal{O}_{44}$ and $\mathcal{O}_{12}$ has been described in detail in our study for the quenched case [2]. The evolution from initially large L /low μ to small L /high μ is obtained by applying the so called step scaling function (SSF) in the Schrödinger functional scheme. Once the perturbative regime is safely reached one continues the evolution in perturbation theory computing the scale and scheme independent RGI matrix element. The connection with experimental results can be obtained by making the adequate perturbative evolution of the RGI matrix element in the $\overline{\text{MS}}$ scheme.

The renormalisation conditions for the local $\mathcal{O}_{44}$ and $\mathcal{O}_{12}$ operators are given by

$$\mathcal{O}_R(\mu) = Z_{\mathcal{O}}^{-1}(a\mu)\mathcal{O}(a), \quad \mathcal{O}_R(\mu = L^{-1}) = \mathcal{O}^{(0)}.$$

The correlators to compute the Z factor are

$$f_{\mathcal{O}}(x_0/L) = -\frac{1}{2} \sum_{\mathbf{x}, \mathbf{y}, \mathbf{z}} \langle \mathcal{O}(\mathbf{x}) \mathbb{S}_q(\mathbf{y}, \mathbf{z}) \rangle$$

$$f_1 = -\frac{1}{2} \sum_{\mathbf{u}, \mathbf{v}, \mathbf{y}, \mathbf{z}} \langle \mathbb{S}'_q(\mathbf{u}, \mathbf{v}) \mathbb{S}_q(\mathbf{y}, \mathbf{z}) \rangle$$

where $\mathbb{S}_q$ and $\mathbb{S}'_q$ are suitable quark sources to probe the operators $\mathcal{O}$. With this definition the renormalisation constants are obtained by

$$Z(a/L, \mu) = c \frac{f_{\mathcal{O}}(x_0/L)}{\sqrt{f_1}}; \quad c = \frac{\sqrt{f_1^{(0)}}}{f_{\mathcal{O}}^{(0)}(x_0/L)}.$$

In order to map the L dependence recursively we use the SSF, rigorously defined on the lattice by

$$\sigma_{Z_{\mathcal{O}}} = \lim_{a \rightarrow 0} \Sigma_{Z_{\mathcal{O}}}(u, a/L)$$

$$\Sigma_{Z_{\mathcal{O}}}(u, a/L) = \frac{Z_{\mathcal{O}}(u, 2L/a)}{Z_{\mathcal{O}}(u, L/a)}, \quad u = \bar{g}_{SF}^2(L).$$

The values of β corresponding to a fixed running coupling for $N_f = 2$ dynamical $O(a)$ -improved




Figure 3. Continuum step scaling function $\sigma_{Z_{44}}$

Wilson fermions are available in [4,5]. We have computed the SSF at six values of the renormalised coupling $\bar{g}_{SF}^2(L) = 0.98$ to 3.33 for the lattice sizes $L/a = 6, 8, 12$, which includes the determination of Z factors on the corresponding L/a and $2L/a$ lattices. The continuum limit has been obtained by a weighted average of several independent runs on the larger $L/a = 8, 12$ (and $2L/a$) lattices using the HMC with one or two pseudo-fermion fields [7]. In fig. 3 we show the $N_f = 2$ continuum running of $\sigma_{Z_{44}}$ in comparison with the previously obtained quenched evolution [2]. In addition we plot the universal 1-loop (dotted) and 2-loop order ($N_f = 0$, dashed) results of perturbation theory. The corresponding plot for the running of $\sigma_{Z_{12}}$ shows a very similar behaviour.

Complemented by a computation of the pion or nucleon matrix element for $N_f = 2$ dynamical fermions, the SSF allows to obtain the RGI matrix element completely non-perturbatively. The whole strategy is summarised by the following formula:

$$\langle \mathcal{O} \rangle_{RGI} = \lim_{a \rightarrow 0} \frac{\langle \mathcal{O} \rangle(a)}{Z_{\mathcal{O}}(a, \mu_0)} \times \sigma_{Z_{\mathcal{O}}}(\mu/\mu_0, \bar{g}^2(\mu)) \mathcal{F}_{SF}(\bar{g}^2(\mu))$$

where we use the $n = 6$ SSF computed at μ_n

$$\sigma\left(\frac{\mu}{\mu_0}, \bar{g}^2(\mu)\right) = \sigma\left(\frac{\mu_1}{\mu_0}, \bar{g}^2(\mu_1)\right) \cdots \sigma\left(\frac{\mu_n}{\mu_{n-1}}, \bar{g}^2(\mu_n)\right)$$

to jump from the non-perturbative scale μ_0 to the

perturbative (ultraviolet) scale μ . At this point one can try to do the perturbative matching using

$$\mathcal{F}_{SF}(\bar{g}^2(\mu)) = [\bar{g}^2(\mu)]^{-\frac{\gamma_0}{2b_0}} \times \exp\left\{-\int_0^{\bar{g}(\mu)} dx \left[\frac{\gamma(x)}{\beta(x)} - \frac{\gamma_0}{2b_0}\right]\right\}$$

computed with 3-loop β and 2-loop γ functions. If the perturbative matching has been successful the quantity

$$\sigma_{INV}^{UV}(\mu_0) = \sigma(\mu/\mu_0, \bar{g}^2(\mu)) \mathcal{F}_{SF}(\bar{g}^2(\mu))$$

should be independent from the ultraviolet scale μ . Indeed we found in the quenched case that the last steps give a very nice plateau [2]. The determination of σ_{INV}^{UV} for $N_f = 2$ dynamical fermions requires a new computation of the 2-loop anomalous dimension γ_1^{SF} in the Schrödinger functional scheme, which is currently work in progress.

ACKNOWLEDGEMENTS

The authors thank F. Knechtli, M. Della Morte and J. Rolf for the generation of dynamical configurations within the running mass project [5] and NIC/DESY Zeuthen for the computer resources.

REFERENCES

1. ZeRo Collaboration: M. Guagnelli et al., hep-lat/0405027
2. ZeRo Collaboration: M. Guagnelli et al., NPB 664 (2003) 276; A. Shindler et al., NPB (PS) 129 (2004) 278
3. ZeRo Collaboration: M. Guagnelli et al., PLB 597 (2004) 216; I. Wetzorke et al., NPB (PS) 129 (2004) 281
4. ALPHA Collaboration: A. Bode et al., PLB 515 (2001) 49; M. Della Morte et al., NPB (PS) 119 (2003) 439
5. ALPHA Collaboration: M. Della Morte et al., CPC 156 (2003) 62; F. Knechtli et al., NPB (PS) 119 (2003) 320; NPB (PS) 129 (2004) 814
6. M. Lüscher et al., NPB 384 (1992) 168; S. Sint, NPB 421 (1994) 135
7. M. Hasenbusch, K. Jansen, NPB 659 (2003) 299